

The University of Reading

Approximate iterative methods for variational data assimilation

A.S. Lawless¹, S. Gratton² and N.K. Nichols¹

NUMERICAL ANALYSIS REPORT 3/04

¹*Department of Mathematics
The University of Reading
Whiteknights, PO Box 220
Reading
Berkshire RG6 6AX*

²*CERFACS
42 Avenue Gustave Coriolis
31057 Toulouse CEDEX
France*

Paper presented at ICFD conference on Numerical Methods for Fluid Dynamics, Oxford, UK,
29 March - 1 April 2004.

Department of Mathematics

with $x \in \mathbb{R}^n$ [3]. We can write the 4D-Var objective function (1) in this form by putting $f(x) = C^{-1/2} \hat{d}$, where

$$\hat{d}(x_0) = \begin{pmatrix} 0 \\ x_0 - x^b \\ H_0[x_0] - y_0^o \\ \vdots \\ H_n[x_n] - y_n^o \end{pmatrix}; \quad C^{-1} = \begin{pmatrix} B_0^{-1} & 0 \\ 0 & R^{-1} \end{pmatrix} \quad \text{with} \quad R = \text{diag}(R_i);$$




Figure 1: Comparison of convergence of (a) objective function and (b) its gradient for a constant convergence criterion (solid line) and a variable criterion (dashed line).

with $D_t \bar{h} = \partial_t \bar{h} + u \partial_x \bar{h}$. In these equations $\bar{h} = \bar{h}(x)$ is the height of the bottom orography, u is the velocity of the fluid and $\bar{A} = gh$ is the geopotential, where g is the gravitational constant and $h > 0$ the depth of the fluid above the orography. The problem is defined on the domain $x \in [0; L]$, with periodic boundary conditions, and we let $t \in [0; T]$.

The model is discretized using a two-time-level semi-implicit semi-Lagrangian scheme. Further details of the numerics can be found in [6]. We use a periodic domain of 1000 grid points, with a spacing $\Delta x = 0.01$ m, so that $x \in [0 \text{ m}; 10 \text{ m}]$. The model time step is taken to be 9.2×10^{-3} s and we consider an assimilation over a window of 100 time steps.

In order to test the assimilation we run ‘identical twin’ experiments, in which observations are generated from a run of the model defined to be the truth. These observations are then assimilated using 4D-Var, starting from an incorrect prior estimate. The inner minimization is performed using a conjugate gradient method and is considered to

Again it is possible to provide a theoretical understanding of these results by an analysis of the (PGN) method. The method can be considered as a way of solving the approximate normal equations

$$\tilde{\mathbf{J}}(\tilde{\mathbf{x}}^*)^T \mathbf{f}(\tilde{\mathbf{x}}^*) = 0; \quad (14)$$

Then based on the work of [7] and [8] we can prove the following theorem:

Theorem 2 *Let the first derivative of $\tilde{\mathbf{J}}(\mathbf{x})^T \mathbf{f}(\tilde{\mathbf{x}})$ be written*

$$\tilde{\mathbf{J}}(\mathbf{x})^T \mathbf{J}(\mathbf{x}) + \tilde{\mathbf{Q}}(\mathbf{x}); \quad (15)$$

where $\tilde{\mathbf{Q}}(\mathbf{x})$ represents second order terms arising from the derivative of $\tilde{\mathbf{J}}(\mathbf{x})$. Suppose that on each iteration k

- [4] Ortega JM, Rheinboldt WC. *Iterative Solution of Nonlinear Equations in Several Variables*