

Department of Mathematics and Statistics

Preprint MPS-2012-11

31 May 2012

Superfast non-linear diffusion in porous media: new regimes, universal model and power laws

by

A.V. Lukyanov, M.M. Sushchikh, M.J. Baines and
T.G. Theofanous



Superfast non-linear diffusion in porous media: new regimes, universal model and power laws.

A.V. Lukyanov¹, M.M. Sushchikh², M.J. Baines¹, T.G. Theofanous²

Department of Mathematics and Statistics, University of Reading, Reading RG6 6AX, U.K.¹

Center for Risk Studies and Safety, University of California at Santa Barbara, Santa Barbara, CA, USA.²

Abstract

The migration of liquids driven by capillary forces in passage networks of porous media, such as sand, has been commonly considered at saturation levels relevant to pore dimensions. In this letter we reveal a very-low-saturation regime where the relevant length scales of the transport mechanics are defined by the grain roughness

pr

the granular medium we use Standard Ottawa Sand (EMD Chemicals, Inc., p/n SX0075); it has an average grain size of $250\text{ }\mu\text{m}$, porosity of 30% and surface roughness in the range $0.25 < \Delta R_0 < 3\text{ }\mu$




Figure 1: Illustration of isolated bridges at low levels of saturations. (a) MicroXCT image, typical from our experiments. (b) 3D image reconstruction of MicroXCT data. The liquid within the roughness of the sand grains,

